

# PHENIX results on multiparticle correlations in small systems


Ron Belmont  
University of Colorado Boulder  
For the PHENIX Collaboration

Quark Matter XXVI  
Chicago, IL  
7 February 2017




University of Colorado  
Boulder

# Testing hydro by controlling system size




J.D. Orjuela Koop et al  
Phys. Rev. C 93, 044910 (2016)



- Standard picture for A+A:  
QGP in hydro evolution
- What about small systems?  
And lower energies?
- Use collision species and  
energy to control system size,  
test limits of hydro applicability

# Multiparticle correlations in small systems

CMS, Phys. Lett. B 765 (2017) 193-220



- Multiparticle correlations: a strong case for collectivity in small systems
- Gaussian fluctuations:

$$v_2\{2\} = \sqrt{v_2^2 + \sigma^2 + \delta}$$

$\delta$  non-flow,  $\sigma$  variance

$$v_2\{2, |\Delta\eta| > 2\} = \sqrt{v_2^2 + \sigma^2}$$

eta gap removes some non-flow


$$v_2\{4\} = v_2\{6\} = v_2\{8\} = \sqrt{v_2^2 - \sigma^2}$$

higher orders remove non-flow


- Can multiparticle correlations be measured in small systems at RHIC?

# 2016 d+Au beam energy scan in PHENIX

200 GeV




62 GeV




d+Au collision energy	total events analyzed	central events analyzed
200 GeV	636 million	585 million
62.4 GeV	131 million	76 million
39 GeV	137 million	49 million
19.6 GeV	15 million	3 million


# The PHENIX forward vertex detector



- FVTX: forward vertex detector
  - silicon strip technology
- Very precise vertex/DCA determination
- No momentum determination,  
 $p_T$  dependent efficiency —  
measured  $v_2$  roughly 18%  
higher than true



# A Multi-Phase Transport model



## AMPT basic features

Initial conditions	MC Glauber
Particle production	String melting
Pre-equilibrium	None
Expansion	Parton scattering (tunable)
Hadronization	Spatial coalescence
Final stage	Hadron cascade (tunable)

- AMPT has significant success in describing flow-like signatures (for low  $p_T$  and  $p_T$ -integrated)
- AMPT produces final state particles over the full available phasespace —possible to perform exact same analysis on data and model

## Experimental method and details

### Definition of Q-vectors

$$Q_{n,x} = \sum_{i=1}^M \cos n\phi_i = \Re e Q_n, \quad Q_{n,y} = \sum_{i=1}^M \sin n\phi_i = \Im m Q_n$$

### Calculation of correlators

$$\langle 2 \rangle = \frac{Q_n Q_n^* - M}{M(M-1)}$$

$$\langle 4 \rangle = \frac{|Q_n|^4 + |Q_{2n}|^2 - 2\Re e [Q_{2n} Q_n^* Q_n^*]}{M(M-1)(M-2)(M-3)} - 2 \frac{2(M-2)|Q_n|^2 - M(M-3)}{M(M-1)(M-2)(M-3)}.$$

### Calculation of cumulants

$$c_n\{2\} = \langle\langle 2 \rangle\rangle$$

$$c_n\{4\} = \langle\langle 4 \rangle\rangle - 2\langle\langle 2 \rangle\rangle^2$$


### Determination of harmonic coefficients

$$v_n\{2\} = \sqrt{c_n\{2\}}$$


$$v_n\{4\} = \sqrt[4]{-c_n\{4\}}$$

# Components and cumulants in p+Au and d+Au at 200 GeV

d+Au




p+Au




- Real  $v_2\{4\}$  in d+Au, imaginary  $v_2\{4\}$  in p+Au
- Fluctuations could dominate in the p+Au ( $v_2\{4\} = \sqrt{v_2^2 - \sigma^2}$ )

# Components and cumulants in p+Au and d+Au in AMPT

d+Au



p+Au



- AMPT similar to data—real  $v_2\{4\}$  in d+Au, imaginary  $v_2\{4\}$  in p+Au
- Fluctuations could dominate in the p+Au ( $v_2\{4\} = \sqrt{v_2^2 - \sigma^2}$ )


## $v_2\{2\}$ and $v_2\{4\}$ in the d+Au beam energy scan

200 GeV

62.4 GeV


39 GeV

19.6 GeV



- $v_2\{2\}$  relatively constant with  $N_{\text{tracks}}^{\text{FVTX}}$  and collision energy

## $v_2\{2\}$ and $v_2\{4\}$ in the d+Au beam energy scan



- $v_2\{2\}$  relatively constant with  $N_{\text{tracks}}^{\text{FVTX}}$  and collision energy
- **Observation of real  $v_2\{4\}$  in d+Au at all energies!!!**
- **Strong evidence for collectivity**


# $v_2\{2\}$ and $v_2\{4\}$ in the d+Au beam energy scan

200 GeV

62.4 GeV


39 GeV

19.6 GeV



79% confidence level that  
 $v_2\{4\}$  is real for  $10 < N_{\text{trk}} < 20$


- Select  $10 < N_{\text{tracks}}^{\text{FVTX}} < 30$ , integrate
- Trend of  $v_2\{2\}$  and  $v_2\{4\}$  merging as  $\sqrt{s_{\text{NN}}}$  is lowered
- AMPT sees the same trend



The story so far:


- Real  $v_2\{4\}$  in d+Au collisions at all energies!
- Imaginary  $v_2\{4\}$  in p+Au at 200 GeV, maybe fluctuations dominate?  
Good reason to believe collectivity/flow in p+Au as well
- Is real-valued  $v_2\{4\}$  really a good measure for collectivity?
- Good news: we can turn the knobs in AMPT to see if we can draw a clear connection between  $v_2\{4\}$  and initial geometry in d+Au

## AMPT with no scattering



- Turn off scattering in AMPT—remove all correlations with initial geometry  
 $\sigma_{parton} = 0$  and  $\sigma_{hadron} = 0$
- Participant plane  $v_2$  goes to zero
- Other sources of correlation remain—non-flow

# AMPT with no scattering



- Turn off scattering in AMPT—remove all correlations with initial geometry
- Components show different trend but are still non-zero
- But  $v_2\{4\}$  goes from real to  $\sim$ zero—connection between real  $v_2\{4\}$  and geometry in d+Au


The story so far:

- Real  $v_2\{4\}$  in d+Au collisions at all energies!
- Clear connection between real  $v_2\{4\}$  and initial geometry—strong evidence for collective behavior

What about the non-flow?


- We've shown  $v_2\{2\}$  but potentially significant non-flow
- We assume  $v_2\{4\}$  removes all the non-flow, but are we sure?
- Try to apply an eta gap on the 2-particle ( $v_2\{2, |\Delta\eta| > 2\}$ ) to get a better handle on non-flow

## How to apply an eta gap in the FVTX?



- $v_2\{2\}$  and  $v_2\{4\}$ —use tracks anywhere in the FVTX

## How to apply an eta gap in the FVTX?



- $v_2\{2\}$  and  $v_2\{4\}$ —use tracks anywhere in the FVTX
- $v_2\{2, |\Delta\eta| > 2\}$ —require one track in south (backward rapidity) and one in north (forward)


Can we apply an eta gap to get a better handle on the non-flow?

200 GeV

62.4 GeV


39 GeV

19.6 GeV



- $v_2\{2\}$  and  $v_2\{4\}$  vs  $N_{\text{tracks}}^{FVTX}$ , all tracks anywhere in FVTX

Can we apply an eta gap to get a better handle on the non-flow?




- $v_2\{2\}$  and  $v_2\{4\}$  vs  $N_{\text{tracks}}^{\text{FVTX}}$ , all tracks anywhere in FVTX
- $v_2\{2, |\Delta\eta| > 2\}$  vs  $N_{\text{tracks}}^{\text{FVTX}}$ , one track backward, the other forward

$$v_2\{2, |\Delta\eta| > 2\} = \sqrt{v_2^2 + \sigma^2} \quad v_2\{2\} = \sqrt{v_2^2 + \sigma^2 + \delta}$$

$$v_2\{4\} = \sqrt{v_2^2 - \sigma^2}$$



- Hard to understand this result based on fluctuations
- The eta gap reduces the non-flow, but what else does it do?

# What can AMPT tell us about asymmetric collisions?





- Asymmetric collision systems have:
  - asymmetric  $dN_{ch}/d\eta$
  - asymmetric  $v_2$  vs  $\eta$
- The FVTX combined is weighted by  $dN_{ch}/d\eta$  towards backward rapidity, where  $v_2$  is also higher—the effect is more pronounced at lower energies
- The FVTX two subevent is equally weighted between forward and back:
$$\sqrt{v_2^B v_2^F}$$

# What can AMPT tell us about asymmetric collisions?




- Asymmetric collision systems have:
  - asymmetric  $dN_{ch}/d\eta$
  - asymmetric  $v_2$  vs  $\eta$
- The FVTX combined is weighted by  $dN_{ch}/d\eta$  towards backward rapidity, where  $v_2$  is also higher—the effect is more pronounced at lower energies
- The FVTX two subevent is equally weighted between forward and back:  
$$\sqrt{v_2^B v_2^F}$$

# What can AMPT tell us about asymmetric collisions?




- Asymmetric collision systems have:
  - asymmetric  $dN_{ch}/d\eta$
  - asymmetric  $v_2$  vs  $\eta$
- The FVTX combined is weighted by  $dN_{ch}/d\eta$  towards backward rapidity, where  $v_2$  is also higher—the effect is more pronounced at lower energies
- The FVTX two subevent is equally weighted between forward and back:
$$\sqrt{v_2^B v_2^F}$$

# Understanding $v_2\{2\}$ , $v_2\{4\}$ , and $v_2\{2, |\Delta\eta| > 2\}$



- $v_2\{2\}$  and  $v_2\{4\}$  vs  $N_{\text{tracks}}^{\text{FVTX}}$ —weighted average of  $v_2^B$  and  $v_2^F$
- $v_2\{2, |\Delta\eta| > 2\}$  vs  $N_{\text{tracks}}^{\text{FVTX}}$ —fixed, equal weighting  $\sqrt{v_2^B v_2^F}$
- $dN_{ch}/d\eta$  and  $v_2$  vs  $\eta$  alone may explain these results

# Understanding $v_2\{2\}$ , $v_2\{4\}$ , and $v_2\{2, |\Delta\eta| > 2\}$



- $v_2\{2\}$  and  $v_2\{4\}$  vs  $N_{\text{tracks}}^{\text{FVTX}}$ —weighted average of  $v_2^B$  and  $v_2^F$
- $v_2\{2, |\Delta\eta| > 2\}$  vs  $N_{\text{tracks}}^{\text{FVTX}}$ —fixed, equal weighting  $\sqrt{v_2^B v_2^F}$
- $dN_{ch}/d\eta$  and  $v_2$  vs  $\eta$  alone may explain these results
- There may be additional effects like event plane decorrelation, e.g.  
 $v_2\{2, |\Delta\eta| > 2\} = \sqrt{v_2^B v_2^F \cos(2(\psi_2^B - \psi_2^F))}$

- Observation of real-valued  $v_2\{4\}$  in d+Au collisions at 200, 62.4, 39, and 19.6 GeV from the 2016 d+Au beam energy scan
  - Connection between real-valued  $v_2\{4\}$  and initial geometry established
  - Strong evidence for collectivity
- $v_2\{4\}$  observed to be everywhere imaginary in p+Au collisions at 200 GeV
  - Could be dominance of fluctuations ( $\sigma/v_2 > 1$ )
  - Good reason to believe collectivity exists in p+Au (many PHENIX measurements on the topic)
  - Contrast with p+Pb at 5.02 TeV—what happens in between RHIC and LHC energies?
- The trend of  $v_2\{2\}$ ,  $v_2\{2, |\Delta\eta| > 2\}$ , and  $v_2\{4\}$  with collision energy is consistent with expectations based on  $dN_{ch}/d\eta$  and  $v_2$  vs  $\eta$ 
  - AMPT shows similar trends
  - Additional effects may be at play—event plane decorrelation?
- Additional measurements are possible and potentially very valuable
  - We plan to explore  $v_2\{6\}$  in d+Au
  - It may be possible to look at  $v_3\{4\}$  in  ${}^3\text{He}+\text{Au}$

### Additional Material

Утро в сосновом лесу




# Утро в сосновом лесу



Почему нам  
нужно заниматься  
физикой?


Почему нет?

# Results




- components, d+Au lower energies, data

# Results




- components, d+Au lower energies, AMPT

# Results




- components, d+Au lower energies, AMPT no scattering

# Results




- $v_2$  in d+Au at all energies

# Results




- $v_2$  in d+Au at all energies

# Results




- $v_2$  in d+Au at all energies

# Results



- $v_2$  in d+Au at all energies

# Results



- $d+\text{Au}$  at 200 GeV on the left,  $p+\text{Au}$  at 200 GeV on right
- $v_2\{2\}$  and  $v_2\{2, |\Delta\eta| > 2\}$  lower for  $p+\text{Au}$ , as expected from geometry